Examination October 2020

B.Sc. S.Y (Sem-III)

2238 Fishery Science Paper-VIII (Culture Fisheries -I)

Time: One Hour Max. Marks: 25

instruction

solve any 25 questions.

1 Fish farming aims at the cultiva	ation of varieties of fishes in ponds.		
(A)Regular	(B)Selected	(C)Normal	(D)Marketed
. , -	pends upon the and the management.	(3)/16/11/61	(D)Markotoa
(A)planning	(B)selection	(C)environment	(D)Manure
3 Selection of suitable for c		(3)3	(2)
(A)Fishes	(B)Water	(C)Sites	(D)Soil
()	ideration in the selection of the site.	(O)Olics	(0)0011
(A)Soil	(B)Topology	(C)Water supply	(D)Stocks
()	h regard to the fertility of fish ponds.	(C)water suppry	(D)Stocks
(A)Topology	(B)Area	(C)Soil	(D)Water
6 A good soil would result in a		(0)3011	(D)vvater
(A)Fish	(B)fertile	(C)Manure	(D)pH
7 The pH of the soil should be in		(C)Manure	(D)pri
(A)6.2-9.0	(B)7.2-9.0	(C)9.0-7.2	(D)4.5-9.0
` '		(0)9.0-7.2	(D)4.3-9.0
(A)Reservoirs	e perhaps the best sources of water.	(C)Panda	(D)Poin
9 Pond depth should be about	(B)Lakes	(C)Ponds	(D)Rain
•		(C)1 5	(D)2 F
(A)1	(B)2	(C)1.5	(D)2.5
10 The size of the pond depends		(C)Stacks	(D)C:
(A)Farm	(B)Area	(C)Stocks	(D)Size
11 These ponds are also known a		(O)-lu- u d-	(D)0
(A)Ponds	(B)flood	(C)sky ponds	(D)Seasonal ponds
12 Hatching pits is also called as	•	(0)	(D)
(A)Rearing	(B)Nursery	(C)spawning	(D)egg
13 Size of hatching tank is fee		(0)01010	(D) 4+0+0
(A)8*4*2	(B)4*2*4	(C)8*2*2	(D)4*8*2
14 The made of coarse cloth			
	or the mosquito curtain cloth.	(0)	(=) II
(A)Net	(В)Нарра	(C)Jal	(D)pit
(A)Net 15 Nursery ponds are also called	(B)Happa as ponds		
(A)Net 15 Nursery ponds are also called (A)Rearing	(B)Happa as ponds (B)transportation	(C)Jal (C)transplantation	(D)pit (D)stocking
(A)Net 15 Nursery ponds are also called (A)Rearing 16 Nursery ponds are constructed	(B)Happa as ponds (B)transportation d of feet	(C)transplantation	(D)stocking
(A)Net 15 Nursery ponds are also called (A)Rearing 16 Nursery ponds are constructed (A)50*50*4	(B)Happa as ponds (B)transportation d of feet (B)40*40*4	(C)transplantation (C)30*40*2	
(A)Net 15 Nursery ponds are also called (A)Rearing 16 Nursery ponds are constructed (A)50*50*4 17 Young fry about days old a	(B)Happa as ponds (B)transportation d of feet (B)40*40*4 are transferred from spawning ponds to nu	(C)transplantation (C)30*40*2 rseries.	(D)stocking (D)40*50*4
(A)Net 15 Nursery ponds are also called (A)Rearing 16 Nursery ponds are constructed (A)50*50*4 17 Young fry about days old a (A)4 to 5	(B)Happa as ponds (B)transportation d of feet (B)40*40*4 are transferred from spawning ponds to nu (B)3 to 5	(C)transplantation (C)30*40*2	(D)stocking
(A)Net 15 Nursery ponds are also called (A)Rearing 16 Nursery ponds are constructed (A)50*50*4 17 Young fry about days old a (A)4 to 5 18 Stocking ponds size may be	(B)Happa as ponds (B)transportation d of feet (B)40*40*4 are transferred from spawning ponds to nu (B)3 to 5 feet.	(C)transplantation (C)30*40*2 rseries. (C)5 to 6	(D)stocking (D)40*50*4 (D)4 to 6
(A)Net 15 Nursery ponds are also called (A)Rearing 16 Nursery ponds are constructed (A)50*50*4 17 Young fry about days old a (A)4 to 5 18 Stocking ponds size may be (A)50*100*6	(B)Happa as ponds (B)transportation d of feet (B)40*40*4 are transferred from spawning ponds to nu (B)3 to 5 feet. (B)300*88*6	(C)transplantation (C)30*40*2 rseries. (C)5 to 6 (C)300*100*5	(D)stocking (D)40*50*4
(A)Net 15 Nursery ponds are also called (A)Rearing 16 Nursery ponds are constructed (A)50*50*4 17 Young fry about days old a (A)4 to 5 18 Stocking ponds size may be (A)50*100*6 19 Ponds is one of the key fac	(B)Happa as ponds (B)transportation d of feet (B)40*40*4 are transferred from spawning ponds to nu (B)3 to 5 feet. (B)300*88*6 ctor in increasing the maximum carrying care	(C)transplantation (C)30*40*2 rseries. (C)5 to 6 (C)300*100*5 apacity.	(D)stocking (D)40*50*4 (D)4 to 6 (D)50*50*6
(A)Net 15 Nursery ponds are also called (A)Rearing 16 Nursery ponds are constructed (A)50*50*4 17 Young fry about days old a (A)4 to 5 18 Stocking ponds size may be (A)50*100*6 19 Ponds is one of the key face (A)Culture	(B)Happa as ponds (B)transportation d of feet (B)40*40*4 are transferred from spawning ponds to nu (B)3 to 5 feet. (B)300*88*6 ctor in increasing the maximum carrying co	(C)transplantation (C)30*40*2 rseries. (C)5 to 6 (C)300*100*5	(D)stocking (D)40*50*4 (D)4 to 6
(A)Net 15 Nursery ponds are also called (A)Rearing 16 Nursery ponds are constructed (A)50*50*4 17 Young fry about days old a (A)4 to 5 18 Stocking ponds size may be (A)50*100*6 19 Ponds is one of the key far (A)Culture 20 has an antiparasitic action.	(B)Happa as ponds (B)transportation d of feet (B)40*40*4 are transferred from spawning ponds to nu (B)3 to 5 feet. (B)300*88*6 ctor in increasing the maximum carrying carrying carrying	(C)transplantation (C)30*40*2 rseries. (C)5 to 6 (C)300*100*5 apacity. (C)fertilization	(D)stocking (D)40*50*4 (D)4 to 6 (D)50*50*6 (D)manure
(A)Net 15 Nursery ponds are also called (A)Rearing 16 Nursery ponds are constructed (A)50*50*4 17 Young fry about days old a (A)4 to 5 18 Stocking ponds size may be (A)50*100*6 19 Ponds is one of the key factory (A)Culture 20 has an antiparasitic action. (A)Fertility	(B)Happa as ponds (B)transportation d of feet (B)40*40*4 are transferred from spawning ponds to nu (B)3 to 5 feet. (B)300*88*6 ctor in increasing the maximum carrying carrying (B)rearing (B)lining	(C)transplantation (C)30*40*2 rseries. (C)5 to 6 (C)300*100*5 apacity.	(D)stocking (D)40*50*4 (D)4 to 6 (D)50*50*6
(A)Net 15 Nursery ponds are also called (A)Rearing 16 Nursery ponds are constructed (A)50*50*4 17 Young fry about days old a (A)4 to 5 18 Stocking ponds size may be (A)50*100*6 19 Ponds is one of the key far (A)Culture 20 has an antiparasitic action. (A)Fertility 21 Overgrowth of phytoplankton research	(B)Happa as ponds (B)transportation d of feet (B)40*40*4 are transferred from spawning ponds to nu (B)3 to 5 feet. (B)300*88*6 ctor in increasing the maximum carrying co (B)rearing (B)lining night cause depletion of	(C)transplantation (C)30*40*2 rseries. (C)5 to 6 (C)300*100*5 apacity. (C)fertilization (C)cow dung	(D)stocking (D)40*50*4 (D)4 to 6 (D)50*50*6 (D)manure (D)cleaning
(A)Net 15 Nursery ponds are also called (A)Rearing 16 Nursery ponds are constructed (A)50*50*4 17 Young fry about days old a (A)4 to 5 18 Stocking ponds size may be (A)50*100*6 19 Ponds is one of the key far (A)Culture 20 has an antiparasitic action. (A)Fertility 21 Overgrowth of phytoplankton r (A)Oxygen	(B)Happa as ponds (B)transportation d of feet (B)40*40*4 are transferred from spawning ponds to nu (B)3 to 5 feet. (B)300*88*6 ctor in increasing the maximum carrying co (B)rearing (B)lining night cause depletion of (B)Carbon	(C)transplantation (C)30*40*2 rseries. (C)5 to 6 (C)300*100*5 apacity. (C)fertilization (C)cow dung (C)Nitrogen	(D)stocking (D)40*50*4 (D)4 to 6 (D)50*50*6 (D)manure
(A)Net 15 Nursery ponds are also called (A)Rearing 16 Nursery ponds are constructed (A)50*50*4 17 Young fry about days old a (A)4 to 5 18 Stocking ponds size may be (A)50*100*6 19 Ponds is one of the key fact (A)Culture 20 has an antiparasitic action. (A)Fertility 21 Overgrowth of phytoplankton r (A)Oxygen 22 The fishes feed upon cultivable	(B)Happa as ponds (B)transportation d of feet (B)40*40*4 are transferred from spawning ponds to nu (B)3 to 5 feet. (B)300*88*6 ctor in increasing the maximum carrying ca (B)rearing (B)lining night cause depletion of (B)Carbon as species of carp hatchlings are called	(C)transplantation (C)30*40*2 rseries. (C)5 to 6 (C)300*100*5 apacity. (C)fertilization (C)cow dung (C)Nitrogen fishes.	(D)stocking (D)40*50*4 (D)4 to 6 (D)50*50*6 (D)manure (D)cleaning (D)carbon dioxide
(A)Net 15 Nursery ponds are also called (A)Rearing 16 Nursery ponds are constructed (A)50*50*4 17 Young fry about days old a (A)4 to 5 18 Stocking ponds size may be (A)50*100*6 19 Ponds is one of the key fac (A)Culture 20 has an antiparasitic action. (A)Fertility 21 Overgrowth of phytoplankton r (A)Oxygen 22 The fishes feed upon cultivable (A)Food	(B)Happa as ponds (B)transportation d of feet (B)40*40*4 are transferred from spawning ponds to nu (B)3 to 5 feet. (B)300*88*6 ctor in increasing the maximum carrying co (B)rearing (B)lining night cause depletion of (B)Carbon e species of carp hatchlings are called (B)predatory	(C)transplantation (C)30*40*2 rseries. (C)5 to 6 (C)300*100*5 apacity. (C)fertilization (C)cow dung (C)Nitrogen	(D)stocking (D)40*50*4 (D)4 to 6 (D)50*50*6 (D)manure (D)cleaning
(A)Net 15 Nursery ponds are also called (A)Rearing 16 Nursery ponds are constructed (A)50*50*4 17 Young fry about days old a (A)4 to 5 18 Stocking ponds size may be (A)50*100*6 19 Ponds is one of the key far (A)Culture 20 has an antiparasitic action. (A)Fertility 21 Overgrowth of phytoplankton r (A)Oxygen 22 The fishes feed upon cultivable (A)Food 23 mixed with water in buck	(B)Happa as ponds (B)transportation d of feet (B)40*40*4 are transferred from spawning ponds to nu (B)3 to 5 feet. (B)300*88*6 ctor in increasing the maximum carrying carrying (B)rearing (B)lining night cause depletion of (B)Carbon as species of carp hatchlings are called (B)predatory set and sprayed over the pond surface.	(C)transplantation (C)30*40*2 rseries. (C)5 to 6 (C)300*100*5 apacity. (C)fertilization (C)cow dung (C)Nitrogen fishes. (C)weed	(D)stocking (D)40*50*4 (D)4 to 6 (D)50*50*6 (D)manure (D)cleaning (D)carbon dioxide (D)none of the above
(A)Net 15 Nursery ponds are also called (A)Rearing 16 Nursery ponds are constructed (A)50*50*4 17 Young fry about days old a (A)4 to 5 18 Stocking ponds size may be (A)50*100*6 19 Ponds is one of the key fac (A)Culture 20 has an antiparasitic action. (A)Fertility 21 Overgrowth of phytoplankton r (A)Oxygen 22 The fishes feed upon cultivable (A)Food 23 mixed with water in buck (A)Croton	(B)Happa as ponds (B)transportation d of feet (B)40*40*4 are transferred from spawning ponds to nu (B)3 to 5 feet. (B)300*88*6 ctor in increasing the maximum carrying co (B)rearing (B)lining night cause depletion of (B)Carbon e species of carp hatchlings are called (B)predatory	(C)transplantation (C)30*40*2 rseries. (C)5 to 6 (C)300*100*5 apacity. (C)fertilization (C)cow dung (C)Nitrogen fishes.	(D)stocking (D)40*50*4 (D)4 to 6 (D)50*50*6 (D)manure (D)cleaning (D)carbon dioxide
(A)Net 15 Nursery ponds are also called (A)Rearing 16 Nursery ponds are constructed (A)50*50*4 17 Young fry about days old a (A)4 to 5 18 Stocking ponds size may be (A)50*100*6 19 Ponds is one of the key fact (A)Culture 20 has an antiparasitic action. (A)Fertility 21 Overgrowth of phytoplankton r (A)Oxygen 22 The fishes feed upon cultivable (A)Food 23 mixed with water in buck (A)Croton 24 Rearing pond size is feet.	(B)Happa as ponds (B)transportation d of feet (B)40*40*4 are transferred from spawning ponds to nu (B)3 to 5 feet. (B)300*88*6 ctor in increasing the maximum carrying carrying (B)rearing (B)lining night cause depletion of (B)Carbon a species of carp hatchlings are called (B)predatory set and sprayed over the pond surface. (B)brassica	(C)transplantation (C)30*40*2 rseries. (C)5 to 6 (C)300*100*5 apacity. (C)fertilization (C)cow dung (C)Nitrogen fishes. (C)weed (C)sugar cane	(D)stocking (D)40*50*4 (D)4 to 6 (D)50*50*6 (D)manure (D)cleaning (D)carbon dioxide (D)none of the above (D)Derris Powder
(A)Net 15 Nursery ponds are also called (A)Rearing 16 Nursery ponds are constructed (A)50*50*4 17 Young fry about days old a (A)4 to 5 18 Stocking ponds size may be (A)50*100*6 19 Ponds is one of the key fac (A)Culture 20 has an antiparasitic action. (A)Fertility 21 Overgrowth of phytoplankton r (A)Oxygen 22 The fishes feed upon cultivable (A)Food 23 mixed with water in buck (A)Croton 24 Rearing pond size is feet. (A)90*30*4	(B)Happa as ponds (B)transportation d of feet (B)40*40*4 are transferred from spawning ponds to nu (B)3 to 5 feet. (B)300*88*6 ctor in increasing the maximum carrying co (B)rearing (B)lining night cause depletion of (B)Carbon e species of carp hatchlings are called (B)predatory ket and sprayed over the pond surface. (B)brassica (B)80*30*4	(C)transplantation (C)30*40*2 rseries. (C)5 to 6 (C)300*100*5 apacity. (C)fertilization (C)cow dung (C)Nitrogen fishes. (C)weed	(D)stocking (D)40*50*4 (D)4 to 6 (D)50*50*6 (D)manure (D)cleaning (D)carbon dioxide (D)none of the above
(A)Net 15 Nursery ponds are also called (A)Rearing 16 Nursery ponds are constructed (A)50*50*4 17 Young fry about days old a (A)4 to 5 18 Stocking ponds size may be (A)50*100*6 19 Ponds is one of the key far (A)Culture 20 has an antiparasitic action. (A)Fertility 21 Overgrowth of phytoplankton r (A)Oxygen 22 The fishes feed upon cultivable (A)Food 23 mixed with water in buck (A)Croton 24 Rearing pond size is feet. (A)90*30*4 25 Fish seed is required for p	(B)Happa as ponds (B)transportation d of feet (B)40*40*4 are transferred from spawning ponds to nu (B)3 to 5 feet. (B)300*88*6 ctor in increasing the maximum carrying ca (B)rearing (B)lining night cause depletion of (B)Carbon a species of carp hatchlings are called (B)predatory ket and sprayed over the pond surface. (B)brassica (B)80*30*4 onds.	(C)transplantation (C)30*40*2 rseries. (C)5 to 6 (C)300*100*5 apacity. (C)fertilization (C)cow dung (C)Nitrogen fishes. (C)weed (C)sugar cane (C)80*20*2	(D)stocking (D)40*50*4 (D)4 to 6 (D)50*50*6 (D)manure (D)cleaning (D)carbon dioxide (D)none of the above (D)Derris Powder (D)70*30*2
(A)Net 15 Nursery ponds are also called (A)Rearing 16 Nursery ponds are constructed (A)50*50*4 17 Young fry about days old a (A)4 to 5 18 Stocking ponds size may be (A)50*100*6 19 Ponds is one of the key fac (A)Culture 20 has an antiparasitic action. (A)Fertility 21 Overgrowth of phytoplankton r (A)Oxygen 22 The fishes feed upon cultivable (A)Food 23 mixed with water in buck (A)Croton 24 Rearing pond size is feet. (A)90*30*4 25 Fish seed is required for p (A)Nursery	(B)Happa as ponds (B)transportation d of feet (B)40*40*4 are transferred from spawning ponds to nu (B)3 to 5 feet. (B)300*88*6 ctor in increasing the maximum carrying ca (B)rearing (B)lining night cause depletion of (B)Carbon a species of carp hatchlings are called (B)predatory set and sprayed over the pond surface. (B)brassica (B)80*30*4 onds. (B)Rearing	(C)transplantation (C)30*40*2 rseries. (C)5 to 6 (C)300*100*5 apacity. (C)fertilization (C)cow dung (C)Nitrogen fishes. (C)weed (C)sugar cane	(D)stocking (D)40*50*4 (D)4 to 6 (D)50*50*6 (D)manure (D)cleaning (D)carbon dioxide (D)none of the above (D)Derris Powder
(A)Net 15 Nursery ponds are also called (A)Rearing 16 Nursery ponds are constructed (A)50*50*4 17 Young fry about days old a (A)4 to 5 18 Stocking ponds size may be (A)50*100*6 19 Ponds is one of the key fact (A)Culture 20 has an antiparasitic action. (A)Fertility 21 Overgrowth of phytoplankton r (A)Oxygen 22 The fishes feed upon cultivable (A)Food 23 mixed with water in buck (A)Croton 24 Rearing pond size is feet. (A)90*30*4 25 Fish seed is required for p (A)Nursery 26 is a prolific breeder and br	(B)Happa as ponds (B)transportation d of feet (B)40*40*4 are transferred from spawning ponds to nu (B)3 to 5 feet. (B)300*88*6 ctor in increasing the maximum carrying co (B)rearing (B)lining night cause depletion of (B)Carbon e species of carp hatchlings are called (B)predatory set and sprayed over the pond surface. (B)brassica (B)80*30*4 onds. (B)Rearing eeds easily in ponds.	(C)transplantation (C)30*40*2 rseries. (C)5 to 6 (C)300*100*5 apacity. (C)fertilization (C)cow dung (C)Nitrogen fishes. (C)weed (C)sugar cane (C)80*20*2 (C)Stocking	(D)stocking (D)40*50*4 (D)4 to 6 (D)50*50*6 (D)manure (D)cleaning (D)carbon dioxide (D)none of the above (D)Derris Powder (D)70*30*2 (D)lake
(A)Net 15 Nursery ponds are also called (A)Rearing 16 Nursery ponds are constructed (A)50*50*4 17 Young fry about days old a (A)4 to 5 18 Stocking ponds size may be (A)50*100*6 19 Ponds is one of the key fac (A)Culture 20 has an antiparasitic action. (A)Fertility 21 Overgrowth of phytoplankton r (A)Oxygen 22 The fishes feed upon cultivable (A)Food 23 mixed with water in buck (A)Croton 24 Rearing pond size is feet. (A)90*30*4 25 Fish seed is required for p (A)Nursery	(B)Happa as ponds (B)transportation d of feet (B)40*40*4 are transferred from spawning ponds to nu (B)3 to 5 feet. (B)300*88*6 ctor in increasing the maximum carrying co (B)rearing (B)lining night cause depletion of (B)Carbon e species of carp hatchlings are called (B)predatory set and sprayed over the pond surface. (B)brassica (B)80*30*4 onds. (B)Rearing eeds easily in ponds. (B)rohu	(C)transplantation (C)30*40*2 rseries. (C)5 to 6 (C)300*100*5 apacity. (C)fertilization (C)cow dung (C)Nitrogen fishes. (C)weed (C)sugar cane (C)80*20*2	(D)stocking (D)40*50*4 (D)4 to 6 (D)50*50*6 (D)manure (D)cleaning (D)carbon dioxide (D)none of the above (D)Derris Powder (D)70*30*2

Examination October 2020

(A)4-5" (D)4-5" (D)4-5"

28 is a floating plant.

(A)Nymphaea (B)Azolla (C)Hydrilla (D)Najas

29 Marginal plants $\ldots\ldots$ grow along the shore line of the ponds.

(A)Typha (B)Lemina (C)pistia (D)Nymphaea

30 Air breathing fish $\ldots\ldots$ is economically important.

(A)Channa (B)Anabes (C)notopterus (D)Clarias